

Catalog No.: 40501

Product Name: PyroRTase™

Description

PyroRTase™, a thermostable M-MLV Reverse Transcriptase, is an RNA-dependent DNA polymerase that synthesizes the complementary cDNA first strand from a single-stranded RNA template with hybridized primer. PyroRTase™ is purified from RNaseH mutant *E. coli* and shows 100% activity at 50°C.

Unit definition

One unit of PyroRTase™ incorporates 1 nmol of dTTP into acid-insoluble material in 10 minutes at 37°C.

Concentration	Package size	Storage
100 U/μL	5 KU/set	-20°C

Storage buffer

20 mM Tris-HCl (pH 7.5), 200 mM NaCl, 0.25 mM EDTA, 0.01% NP-40(v/v), 2.5 mM DTT, 50% glycerol (v/v).

5X RT Buffer

250 mM Tris-HCl (pH 8.3), 15 mM MgCl₂, 375 mM KCl, 50 mM DTT.

Features

- **Lack RNase H activity:** Weak RNaseH activity. High yield of long length cDNA.
- **Thermal stable:** Stable from 50°C to 60°C. Allow to reduce the effect of secondary structure of RNA during cDNA synthesis.
- **Wide temperature range:** Capable to perform from 37°C to 60°C.

- **Strong amplification activity:** Gene mutation enhanced the amplification speed and cDNA. Suitable for cDNA library construction.

Recommended Reaction Conditions:

The first-strand cDNA synthesis

- 1) Add the following reagents to a RNase free PCR tube at room temperature.

Oligo dT ₁₂₋₁₈ (1 μg) or random primer (50-250 ng) or gene-specific primer (2 pmole)	x μL
Total RNA (10 ng - 5 μg) or mRNA (1-500 ng)	x μL
dNTP (10 mM each; Cat. No. 40701)	1 μL
DEPC ddH ₂ O	to 14 μL

- 2) Gently mix and incubate 10 min at 70°C then chill on ice for 2-10 min.
- 3) Centrifuge for few seconds then add the next composition on ice:

5X RT Buffer	4 μL
RNase inhibitor (10 U/μL)	1 μL

- 4) Gently mix and incubate at 50°C for 2 min (for Oligo dT₁₂₋₁₈ or gene-specific primer) or at 25°C for 10 min (for random primer).
- 5) Centrifuge for seconds before adding 1 μL PyroRTase™ (100 U/μL). Incubate at 45°C to 50°C for 50-70 min.
- 6) Inactivate PyroRTase™ at 70°C for 10 min.

For research use only.

Leadgene Biomedical, Inc.

TEL: +886-6-505-2959

Address: Rm. R402, No.12, Nanke 2nd Rd., Xinshi Dist., Tainan City 744, Taiwan

Email: info@leadgene.com.tw

www.leadgene.com.tw